



10/538129  
Rec'd PCT/PTO 06 JUN 2005  
PCT/EP / 13803  
EP03/13803  
INVESTOR IN PEOPLE  
#2

EPO - DG 1

14. 01. 2004

(79)

The Patent Office  
Concept House  
Cardiff Road  
Newport  
South Wales  
NP10 8QQ

REC'D 23 JAN 2004

WIPO PCT

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.

**PRIORITY DOCUMENT**  
SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH  
RULE 17.1(a) OR (b)

Signed

*Stephen Handley*

Dated 3 December 2003

**BEST AVAILABLE COPY**

Patents Form 1/77

Patents Act 1977  
(Rule 16)

The  
Patent  
Office

1/77

Request for grant of a patent

(See the notes on the back of this form. You can also get  
an explanatory leaflet from the Patent Office to help  
you fill in this form)



The Patent Office  
Cardiff Road  
Newport  
Gwent NP9 1RH

1. Your reference

RJE/RSG/PG5044

2. Patent application number

(The Patent Office will fill in his part)

0228540.1

06 DEC 2002

3. Full name, address and postcode of the or of  
each applicant (*underline all surnames*)

Glaxo Group Limited  
Glaxo Wellcome House, Berkeley Avenue,  
Greenford, Middlesex UB6 0NN, Great Britain

Patents ADP number (*if you know it*)

If the applicant is a corporate body, give the  
country/state of its incorporation

United Kingdom

004435 84003

4. Title of the invention

Novel Compositions

5. Name of your agent (*if you have one*)

Corporate Intellectual Property

"Address for service" in the United Kingdom  
to which all correspondence should be sent  
(*including the postcode*)

GlaxoSmithKline  
Corporate Intellectual Property (CN9 25.1)  
980 Great West Road  
BRENTFORD  
Middlesex TW8 9GS

Patents ADP number (*if you know it*)

01960 982003

6. If you are declaring priority from one or more  
earlier patent applications, give the country  
and the date of filing of the or each of  
these earlier applications and (*if you know it*) the  
or each application number

Country      Priority application number      Date of filing  
(*if you know it*)      (*day / month / year*)

7. If this application is divided or otherwise  
derived from an earlier UK application,  
give the number and the filing date of  
the earlier application

Number of earlier application      Date of filing  
(*day / month / year*)

8. Is a statement of inventorship and of right

**Patents Form 1/77**

9. Enter the number of sheets for any of the following items you are filing with this form. Do not count copies of the same document

Continuation sheets of this form  
Description  
Claim(s)  
Abstract  
Drawings

20



10. If you are also filing any of the following, state how many against each item.

**Priority Documents**

Translations of priority documents

Statement of inventorship and right to grant of a patent (*Patents Form 7/77*)

Request for preliminary examination and search (*Patents Form 9/77*)

Request for substantive examination (*Patents Form 10/77*)

Any other documents  
(please specify)

11.

We request the grant of a patent on the basis of this application

Signature

R J Easeman

Date 6-Dec-02

12. Name and daytime telephone number of person to contact in the United Kingdom

R J Easeman 020 80474407

**Warning**

*After an application for a Patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission unless an application has been filed at least six weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.*

**Notes**

- a) If you need help to fill in this form or you have any questions, please contact the Patent Office on 0645 500505
- b) Write your answers in capital letters using black ink or you may type them.
- c) If there is not enough space for all relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be

## Novel Compositions

The present invention relates to compositions comprising DNA attached to an immunostimulatory oligonucleotide (CpG) via a locked nucleic acid oligonucleotide.

5 In particular the present invention provides compositions comprising a plasmid containing a gene encoding a protein of interest, wherein said plasmid may be introduced to a tissue or cell and the gene expressed, complexed to the LNA -CpG.

Plasmid based delivery of genes, particularly for immunisation or gene therapy purposes is known. For example, administration of naked DNA by injection  
10 into mouse muscle is outline by Vical in International Patent Application WO90/11092.

Johnston et al WO 91/07487 describe methods of transferring a gene to vertebrate cells, by the use of microprojectiles that have been coated with a polynucleotide encoding a gene of interest, and accelerating the microparticles such  
15 that the microparticles can penetrate the target cell.

DNA vaccines usually consist of a bacterial plasmid vector into which is inserted a strong viral promoter, the gene of interest which encodes for an antigenic peptide and a polyadenylation/transcriptional termination sequences. The gene of interest may encode a full protein or simply an antigenic peptide sequence relating to  
20 the pathogen, tumour or other agent which is intended to be protected against. The plasmid can be grown in bacteria, such as for example *E.coli* and then isolated and prepared in an appropriate medium, depending upon the intended route of administration, before being administered to the host. Following administration the plasmid is taken up by cells of the host where the encoded peptide is produced. The  
25 plasmid vector will preferably be made without an origin of replication which is functional in eukaryotic cells, in order to prevent plasmid replication in the mammalian host and integration within chromosomal DNA of the animal concerned.

There are a number of advantages of DNA vaccination relative to traditional vaccination techniques. First, it is predicted that because of the proteins which are  
30 encoded by the DNA sequence are synthesised in the host, the structure or conformation of the protein will be similar to the native protein associated with the disease state. It is also likely that DNA vaccination will offer protection against different strains of a virus, by generating cytotoxic T lymphocyte response that

recognise epitopes from conserved proteins. Furthermore, because the plasmids are taken up by the host cells where antigenic protein can be produced, a long-lasting immune response will be elicited. The technology also offers the possibility of combining diverse immunogens into a single preparation to facilitate simultaneous immunisation in relation to a number of disease states.

Helpful background information in relation to DNA vaccination is provided in Donnelly et al "DNA vaccines" Ann. Rev Immunol. 1997 15: 617-648, the disclosure of which is included herein in its entirety by way of reference.

Despite the numerous advantages associated with DNA vaccination relative to traditional vaccination therapies there is nonetheless a desire to develop improvements which will serve to increase the immune response induced by the protein which is encoded by the plasmid DNA administered to an animal. The present invention addresses these issues.

Locked nucleic acid (LNA) is an analogue of RNA or DNA. The term LNA is used to describe both nucleotide monomers, in which the ribose ring is constrained by a methylene linkage between the 2' - oxygen and the 4' - carbon, and also oligonucleotides that contain one or more monomers of locked nucleic acid. The methylene bridge linkage can be through oxygen, (oxy-LNA), sulphur, (thio-LNA) and amine, (amino-LNA). The confirmation restriction increases binding affinity for complementary sequences (Dwayne A. Braasch and David R. Corey, Chemistry and Biology 8 (2001) 1-7). The introduction of LNA monomers into DNA or RNA oligonucleotides increases affinity for complementary DNA or RNA sequences, ie. measured as thermal stability of duplexes, eg. melting temperature, ( $T_m$ ), increases in the range of 3 - 8°C, depending on the actual base, per LNA monomer present in the oligonucleotide., (Dwayne A. Braasch and David R. Corey, Chemistry and Biology 8 (2001) 1-7). Synthesis of LNA is described in International Patent Application No. WO99/14226.

Although triplex formation of LNA oligonucleotides with short double stranded DNA oligonucleotides has been described, (74, 75), no report has yet been published on the properties of LNA oligonucleotides as strand displacement agents in conjunction with large supercoiled plasmid DNA molecules, which in part is the subject matter of the present invention. Indeed, it has been recently suggested that the charged backbone of LNA oligonucleotides would make them less efficient strand

displacement agents than uncharged PNA oligonucleotides, (Dwayne A. Braasch and David R. Corey, Chemistry and Biology 8 (2001) 1-7). The present inventors here provide clear evidence that LNA oligonucleotides are at least as efficient strand displacement agents of supercoiled plasmid DNA as their PNA derived counterparts.

5 We also provide strong evidence that once bound to plasmid DNA, LNA oligonucleotides are more stably attached to plasmid DNA than PNA oligonucleotides and can remain bound when exposed to harsh condition whereas PNA oligonucleotides do not. This is advantageous when considering formulating such DNA/LNA complexes for pharmaceutical administration.

10 Linking peptide and other material (eg. fluorescent labels such as rhodamine) to DNA plasmid by means of a Peptide nucleic acid oligonucleotide is known (US 6165720). These have also been used to transfect cells. Although reports have been made of PNA / DNA / PNA triplexes surviving quite harsh conditions (50), such studies were only performed on short DNA oligonucleotides and not upon large  
15 supercoiled plasmid DNA, where stability of bound PNA to a range of external conditions has not been reported. Moreover, it has been found by the present inventors that such complexes are not sufficiently stable to enable a PNA-coupled fluorophore or peptide to remain attached to plasmid DNA when administered in a pharmaceutical or vaccine formulation, especially for (particle mediated  
20 immunotherapeutic delivery) PMID. Additionally, the inventors have found labelling of plasmid DNA with PNA oligonucleotides to have variable efficiency, poor reproducibility and constraints on reaction conditions in requiring low or no salt and low pH, (<6), for optimal PNA labelling.

The present invention provides a novel LNA- immunostimulatory  
25 oligonucleotide compositions that may be used enhance the immune response stimulated by DNA vaccines. The immunostimulatory oligonucleotides in the present invention are oligonucleotides that comprise at least one unmethylated CG dinucleotide (CpG).

30 In the first aspect of the present invention there is provided a novel composition comprising an oligonucleotide of the following CpG sequence:

5'-tccatgacgttcctgacgtt-3' SEQ ID No. 1.

attached to the following LNA sequence, optionally through one or more nucleotide linker residue:

5'-GGAAGGAAGGAAGG-3', SEQ ID No. 2.

5 Preferably the LNA-CpG sequence is:

5'-tccatgacgttcctgacgttXGGAAGGAAGGAAGG-3' SEQ. ID No. 3

wherein X can be A, G, T or C, but preferably T.

In a second aspect of the present invention there is provided a method of manufacturing a LNA – CpG conjugate comprising co-synthesising the entire sequence. Preferably the oligonucleotide produced by this method is SEQ ID No.3. Conjugates formed by the method of the present invention also form an aspect of the present invention.

Also forming an aspect of the present invention are methods of producing DNA plasmid vaccines by binding these LNA/CpG conjugates to plasmid DNA containing a gene under the control of a promoter such that the gene may be expressed *in vivo*. The LNA conjugate is stable and can be administered *in vivo* with the plasmid DNA allowing co-localisation of the plasmid and the CpG within the cells whilst still retaining the ability of the gene to be expressed. LNA oligonucleotides, advantageously are not subject to degradation by intracellular Dnase enzymes, (Dwayne A. Braasch and David R. Corey, Chemistry and Biology 8 (2001) 1-7).

The LNA conjugate produced by the method of the second aspect of the present invention comprises an oligonucleotide of between 7-25, preferably 10-20, more preferably 11-15 bases at least one of which is a locked nucleic acid preferably at least half, more preferably the entire oligonucleotide is made of LNA bases.

Typically, at least a sequence of at least 13 LNA residues is preferred for optimal stability, when bound to the plasmid DNA. Preferred LNA molecules for use in the second aspect of the present invention are listed in Tables 1. Particularly preferred LNA oligonucleotides are shown in table 1 as LNA 4, or SEQ ID No.2. The LNA oligonucleotide should be free from self-complementary base-pairing sequences for optimal binding to DNA. An alternative embodiment can be envisaged where complementary sequences to further LNA oligonucleotides are present in initial bound LNA oligonucleotides such that an array of LNA oligonucleotide can be bound to a

single LNA complementary site within DNA, formed by LNA : LNA hybridization between LNA oligonucleotides.

In the present invention the LNA is conjugated to a CpG immunostimulatory moiety, so that once the LNA/CpG is associated with the DNA plasmid encoding a gene of interest and administered to a host, the DNA plasmid can express the gene and allow the function of the attached moiety.

LNA and CpG, and direct co-synthesis of the two may be performed by methods as described in Verma, S. & Eckstein F. Annual Review of Biochemistry 1998, 67: 99-134, Modified oligonucleotides: synthesis and strategy for users.

ed. M.J. Gait. Oligonucleotide synthesis: a practical approach- book IRL / Oxford University Press 1990

ed Sudhir Agrawal. Methods in Molecular Biology Vol 20: Protocols for oligonucleotides and analogs, synthesis and properties- book Humana Press, 1993.

Particularly preferred adjuvants for linking to DNA plasmids via the LNA are CpG oligo- and di-nucleotides, (65, 66). The CpG immunostimulatory sequence is often: Purine, Purine, C, G, pyrimidine, pyrimidine; wherein the dinucleotide CG motif is not methylated . The preferred oligonucleotides for use in adjuvants or vaccines of the present invention preferably contain two or more dinucleotide CpG motifs separated by at least three, more preferably at least six or more nucleotides. The oligonucleotides of the present invention are typically deoxynucleotides. In a preferred embodiment the internucleotide in the oligonucleotide is phosphorodithioate, or more preferably a phosphorothioate bond, although phosphodiester and other internucleotide bonds are within the scope of the invention including oligonucleotides with mixed internucleotide linkages. Methods for producing phosphorothioate oligonucleotides or phosphorodithioate are described in US5,666,153, US5,278,302 and WO95/26204.

Examples of preferred oligonucleotides have the following sequences. The sequences preferably contain phosphorothioate modified internucleotide linkages.

OLIGO 1 (SEQ ID NO:4): TCC ATG ACG TTC CTG ACG TT (CpG 1826)

OLIGO 2 (SEQ ID NO:5): TCT CCC AGC GTG CGC CAT (CpG 1758)

OLIGO 3 (SEQ ID NO:6): ACC GAT GAC GTC GCC GGT GAC GGC ACC ACG

OLIGO 4 (SEQ ID NO:1): TCG TCG TTT TGT CGT TTT GTC GTT (CpG 2006)



OLIGO 5 (SEQ ID NO:8): TCC ATG ACG TTC CTG ATG CT (CpG 1668)

Alternative CpG oligonucleotides may comprise the preferred sequences above in that they have inconsequential deletions or additions thereto.

The CpG oligonucleotides utilised in the present invention may be synthesised by any  
5 method known in the art (e.g. EP 468520). Conveniently, such oligonucleotides may be synthesised utilising an automated synthesiser.

The oligonucleotides utilised in the present invention are typically deoxynucleotides. In a preferred embodiment the internucleotide bond in the oligonucleotide is phosphorodithioate, or more preferably phosphorothioate bond, although  
10 phosphodiester are within the scope of the present invention. Oligonucleotide comprising different internucleotide linkages are contemplated, e.g. mixed phosphorothioate phosphodiester. Other internucleotide bonds which stabilise the oligonucleotide may be used.

In the first aspect of the present invention the CpG may be chemically  
15 conjugated to the LNA using any of a range of commercially available cross-linking reagents. The examples described below are by no means exhaustive and include utilisation of the amino, aryl, carboxyl and hydroxyl groups found on peptides or proteins and have been extensively reviewed, (2). Other heterobifunctional cross-linking reagents are available for coupling such reactive groups including  
20 carbodiimide cross-linkers to couple carboxyl groups to amines, eg. 1-ethyl-3-(3-dimethylaminopropyl) - carbodiimide hydrochloride and other cross-linking reagents that couple to sulphhydryl groups, (eg. haloacetyls or pyridyl disulphide), or amino groups, eg. imidoesters or N-hydroxysuccinimide-esters including succinimidyl 4-(N-maleimidomethyl) cyclohexane-1-carboxylate (SMCC) and succinimidyl-4-(p-  
25 maleimidophenyl)-butyrate (SMPB).

One preferred embodiment in either of the first two aspects this invention is to design the linkage of the CpG to the LNA oligonucleotide such that it can be selectively cleaved, (perhaps in order to exert a biological response), from the LNA oligonucleotide and bound plasmid once they have been delivered to a cell. One such  
30 example of this is where the CpG adjuvant, as a phosphorothioate oligonucleotide, is linked to an LNA oligonucleotide by a single DNA phosphoramidate residue, which leaves the 'hybrid' oligonucleotide available for cleavage by cellular phosphodiester enzymes upon delivery to the endosomal compartment of the cell. Cleavage could then

release the CpG adjuvant as a free phosphorothioate oligonucleotide to exert its biological effect.

5 In a preferred embodiment of the present invention the LNA – conjugate is associated with a DNA molecule encoding a gene, said DNA molecule having a sequence complementary to the LNA oligonucleotide. The DNA is preferably in the form of a plasmid and preferably encodes an antigen or therapeutic protein.

The plasmid is preferably without a functional origin of replication in order to prevent plasmid replication in the host to which it is administered. The promoter is preferably a strong viral promoter such as a CMV promoter.

10 The plasmid can be provided with a plurality of LNA complementary binding sequences to enable a plurality of LNA/conjugates to bind. The conjugates may have discrete different functional moities. Thus in one aspect of the invention the plasmid may bind to an LNA linked to a nuclear localisation peptide and an LNA linked to a small molecule adjuvant. Typically the plasmid will be provided with 4 or more  
15 complementary LNA binding sequences preferably 10 to 20 sequences, but up to 100 sequences are possible. Accordingly in one aspect of the invention there is provided a plasmid LNA conjugate complex wherein there is at least four LNA conjugates bound to the plasmid.

In a preferred embodiment the antigen is capable of eliciting an immune  
20 response against a human pathogen, which antigen or antigenic composition is derived from HIV-1, (such as tat, nef, gp120 or gp160, gp40, p24, gag, env, vif, vpr, vpu, rev), human herpes viruses, such as gH, gL gM gB gC gK gE or gD or derivatives thereof or Immediate Early protein such as ICP27 , ICP 47, IC P 4, ICP36 from HSV1 or HSV2, cytomegalovirus, especially Human, (such as gB or derivatives  
25 thereof), Epstein Barr virus (such as gp350 or derivatives thereof), Varicella Zoster Virus (such as gpI, II, III and IE63), or from a hepatitis virus such as hepatitis B virus (for example Hepatitis B Surface antigen or Hepatitis core antigen or pol), hepatitis C virus antigen and hepatitis E virus antigen, or from other viral pathogens, such as paramyxoviruses: Respiratory Syncytial virus (such as F and G proteins or derivatives  
30 thereof), or antigens from parainfluenza virus, measles virus, mumps virus, human papilloma viruses (for example HPV6, 11, 16, 18, eg L1, L2, E1, E2, E3, E4, E5, E6, E7), flaviviruses (e.g. Yellow Fever Virus, Dengue Virus, Tick-borne encephalitis virus, Japanese Encephalitis Virus) or Influenza virus cells, such as HA, NP, NA, or

M proteins, or combinations thereof), or antigens derived from bacterial pathogens such as *Neisseria spp*, including *N. gonorrhea* and *N. meningitidis*, eg, transferrin-binding proteins, lactoferrin binding proteins, PilC, adhesins); *S. pyogenes* (for example M proteins or fragments thereof, C5A protease, *S. agalactiae*, *S. mutans*; *H. ducreyi*; *Moraxella spp*, including *M. catarrhalis*, also known as *Branhamella catarrhalis* (for example high and low molecular weight adhesins and invasins); *Bordetella spp*, including *B. pertussis* (for example pertactin, pertussis toxin or derivatives thereof, filamentous hemagglutinin, adenylate cyclase, fimbriae), *B. parapertussis* and *B. bronchiseptica*; *Mycobacterium spp.*, including *M. tuberculosis* (for example ESAT6, Antigen 85A, -B or -C, MPT 44, MPT59, MPT45, HSP10, HSP65, HSP70, HSP 75, HSP90, PPD 19kDa [Rv3763], PPD 38kDa [Rv0934] ), *M. bovis*, *M. leprae*, *M. avium*, *M. paratuberculosis*, *M. smegmatis*; *Legionella spp*, including *L. pneumophila*; *Escherichia spp*, including enterotoxigenic *E. coli* (for example colonization factors, heat-labile toxin or derivatives thereof, heat-stable toxin or derivatives thereof), enterohemorrhagic *E. coli*, enteropathogenic *E. coli* (for example shiga toxin-like toxin or derivatives thereof); *Vibrio spp*, including *V. cholera* (for example cholera toxin or derivatives thereof); *Shigella spp*, including *S. sonnei*, *S. dysenteriae*, *S. flexnerii*; *Yersinia spp*, including *Y. enterocolitica* (for example a Yop protein) , *Y. pestis*, *Y. pseudotuberculosis*; *Campylobacter spp*, including *C. jejuni* (for example toxins, adhesins and invasins) and *C. coli*; *Salmonella spp*, including *S. typhi*, *S. paratyphi*, *S. choleraesuis*, *S. enteritidis*; *Listeria spp.*, including *L. monocytogenes*; *Helicobacter spp*, including *H. pylori* (for example urease, catalase, vacuolating toxin); *Pseudomonas spp*, including *P. aeruginosa*; *Staphylococcus spp.*, including *S. aureus*, *S. epidermidis*; *Enterococcus spp.*, including *E. faecalis*, *E. faecium*; *Clostridium spp.*, including *C. tetani* (for example tetanus toxin and derivative thereof), *C. botulinum* (for example botulinum toxin and derivative thereof), *C. difficile* (for example clostridium toxins A or B and derivatives thereof); *Bacillus spp.*, including *B. anthracis* (for example botulinum toxin and derivatives thereof); *Corynebacterium spp.*, including *C. diphtheriae* (for example diphtheria toxin and derivatives thereof); *Borrelia spp.*, including *B. burgdorferi* (for example OspA, OspC, DbpA, DbpB), *B. garinii* (for example OspA, OspC, DbpA, DbpB), *B. afzelii* (for example OspA, OspC, DbpA, DbpB), *B. andersonii* (for example OspA, OspC, DbpA, DbpB), *B. hermsii*; *Ehrlichia spp.*,

including *E. equi* and the agent of the Human Granulocytic Ehrlichiosis; *Rickettsia* spp., including *R. rickettsii*; *Chlamydia* spp., including *C. trachomatis* (for example MOMP, heparin-binding proteins), *C. pneumoniae* (for example MOMP, heparin-binding proteins), *C. psittaci*; *Leptospira* spp., including *L. interrogans*; *Treponema* spp., including *T. pallidum* (for example the rare outer membrane proteins), *T. denticola*, *T. hyodysenteriae*; or derived from parasites such as *Plasmodium* spp., including *P. falciparum*; *Toxoplasma* spp., including *T. gondii* (for example SAG2, SAG3, Tg34); *Entamoeba* spp., including *E. histolytica*; *Babesia* spp., including *B. microti*; *Trypanosoma* spp., including *T. cruzi*; *Giardia* spp., including *G. lamblia*; *Leshmania* spp., including *L. major*; *Pneumocystis* spp., including *P. carinii*; *Trichomonas* spp., including *T. vaginalis*; *Schistosoma* spp., including *S. mansoni*, or derived from yeast such as *Candida* spp., including *C. albicans*; *Cryptococcus* spp., including *C. neoformans*.

Other preferred specific antigens for *M. tuberculosis* are for example Rv2557, Rv2558, RPFs: Rv0837c, Rv1884c, Rv2389c, Rv2450, Rv1009, aceA (Rv0467), PstS1, (Rv0932), SodA (Rv3846), Rv2031c 16kDal., Tb Ra12, Tb H9, Tb Ra35, Tb38-1, Erd 14, DPV, MTI, MSL, mTTC2 and hTCC1 (WO 99/51748). Proteins for *M. tuberculosis* also include fusion proteins and variants thereof where at least two, preferably three polypeptides of *M. tuberculosis* are fused into a larger protein. Preferred fusions include Ra12-TbH9-Ra35, Erd14-DPV-MTI, DPV-MTI-MSL, Erd14-DPV-MTI-MSL-mTCC2, Erd14-DPV-MTI-MSL, DPV-MTI-MSL-mTCC2, TbH9-DPV-MTI (WO 99/51748).

Most preferred antigens for *Chlamydia* include for example the High Molecular Weight Protein (HWMP) (WO 99/17741), ORF3 (EP 366 412), and putative membrane proteins (Pmps). Other *Chlamydia* antigens of the vaccine formulation can be selected from the group described in WO 99/28475.

Preferred bacterial vaccines comprise antigens derived from *Streptococcus* spp., including *S. pneumoniae* (PsaA, PspA, streptolysin, choline-binding proteins) and the protein antigen Pneumolysin (Biochem Biophys Acta, 1989, 67, 1007; Rubins et al., Microbial Pathogenesis, 25, 337-342), and mutant detoxified derivatives thereof (WO 90/06951; WO 99/03884). Other preferred bacterial vaccines comprise antigens derived from *Haemophilus* spp., including *H. influenzae* type B (for example PRP and conjugates thereof), non typeable *H. influenzae*, for example OMP26, high molecular

weight adhesins, P5, P6, protein D and lipoprotein D, and fimbrin and fimbrin derived peptides (US 5,843,464) or multiple copy variants or fusion proteins thereof.

The antigens that may be used in the present invention may further comprise antigens derived from parasites that cause Malaria. For example, preferred antigens from *Plasmodia falciparum* include RTS,S and TRAP. RTS is a hybrid protein comprising substantially all the C-terminal portion of the circumsporozoite (CS) protein of *P. falciparum* linked via four amino acids of the preS2 portion of Hepatitis B surface antigen to the surface (S) antigen of hepatitis B virus. Its full structure is disclosed in the International Patent Application No. PCT/EP92/02591, published under Number WO 93/10152 claiming priority from UK patent application No.9124390.7. When expressed in yeast RTS is produced as a lipoprotein particle, and when it is co-expressed with the S antigen from HBV it produces a mixed particle known as RTS,S. TRAP antigens are described in the International Patent Application No. PCT/GB89/00895, published under WO 90/01496. A preferred embodiment of the present invention is a Malaria vaccine wherein the antigenic preparation comprises a combination of the RTS, S and TRAP antigens. Other plasmodia antigens that are likely candidates to be components of a multistage Malaria vaccine are *P. falciparum* MSP1, AMA1, MSP3, EBA, GLURP, RAP1, RAP2, Sequestrin, PfEMP1, Pf332, LSA1, LSA3, STARP, SALSA, PfEXP1, Pfs25, Pfs28, PFS27/25, Pfs16, Pfs48/45, Pfs230 and their analogues in *Plasmodium* spp.

The invention contemplates the use of an anti-tumour antigen and be useful for the immunotherapeutic treatment of cancers. For example, tumour rejection antigens such as those for prostate, breast, colorectal, lung, pancreatic, renal or melanoma cancers. Exemplary antigens include MAGE 1, 3 and MAGE 4 or other MAGE antigens such as disclosed in WO99/40188, PRAME, BAGE, Lage (also known as NY Eos 1) SAGE and HAGE (WO 99/53061) or GAGE (Robbins and Kawakami, 1996, Current Opinions in Immunology 8, pps 628-636; Van den Eynde et al., International Journal of Clinical & Laboratory Research (submitted 1997); Correale et al. (1997), Journal of the National Cancer Institute 89, p293. Indeed these antigens are expressed in a wide range of tumour types such as melanoma, lung carcinoma, sarcoma and bladder carcinoma.

MAGE antigens for use in the present invention may be expressed as a fusion protein with an expression enhancer or an Immunological fusion partner. In

particular, the Mage protein may be fused to Protein D from *Haemophilus influenzae* B. In particular, the fusion partner may comprise the first 1/3 of Protein D. Such constructs are disclosed in Wo99/40188. Other examples of fusion proteins that may contain cancer specific epitopes include *bcr / abl* fusion proteins.

5 In a preferred embodiment prostate antigens are utilised, such as Prostate specific antigen (PSA), PAP, PSCA (PNAS 95(4) 1735-1740 1998), PSMA or antigen known as Prostase.

10 Prostase is a prostate-specific serine protease (trypsin-like), 254 amino acid-long, with a conserved serine protease catalytic triad H-D-S and a amino-terminal pre-peptide sequence, indicating a potential secretory function (P. Nelson, Lu Gan, C. Ferguson, P. Moss, R. Gelinas, L. Hood & K. Wand, "Molecular cloning and characterisation of prostase, an androgen-regulated serine protease with prostate restricted expression, *In Proc. Natl. Acad. Sci. USA* (1999) 96, 3114-3119). A putative glycosylation site has been described. The predicted structure is very similar  
15 to other known serine proteases, showing that the mature polypeptide folds into a single domain. The mature protein is 224 amino acids-long, with one A2 epitope shown to be naturally processed.

Prostase nucleotide sequence and deduced polypeptide sequence and homologs are disclosed in Ferguson, et al. (*Proc. Natl. Acad. Sci. USA* 1999, 96,  
20 3114-3119) and in International Patent Applications No. WO 98/12302 (and also the corresponding granted patent US 5,955,306), WO 98/20117 (and also the corresponding granted patents US 5,840,871 and US 5,786,148) (prostate-specific kallikrein) and WO 00/04149 (P703P).

The present invention provides antigens comprising prostase protein fusions  
25 based on prostase protein and fragments and homologues thereof ("derivatives"). Such derivatives are suitable for use in therapeutic vaccine formulations which are suitable for the treatment of a prostate tumours. Typically the fragment will contain at least 20, preferably 50, more preferably 100 contiguous amino acids as disclosed in the above referenced patent and patent applications.

30 A further preferred prostate antigen is known as P501S, sequence ID no 113 of WO98/37814. Immunogenic fragments and portions encoded by the gene thereof comprising at least 20, preferably 50, more preferably 100 contiguous amino acids as

disclosed in the above referenced patent application, are contemplated. A particular fragment is PS108 (WO 98/50567).

Other prostate specific antigens are known from WO98/37418, and WO/004149. Another is STEAP PNAS 96 14523 14528 7-12 1999.

5 Other tumour associated antigens useful in the context of the present invention include: Plu-1 J Biol. Chem 274 (22) 15633-15645, 1999, HASH-1, HasH-2, Cripto (Salomon et al Bioessays 199, 21 61-70, US patent 5654140) Criptin US patent 5 981 215, ., Additionally, antigens particularly relevant for vaccines in the therapy of cancer also comprise tyrosinase and survivin.

10 The present invention is also useful in combination with breast cancer antigens such as Muc-1, Muc-2, EpCAM, her 2/ Neu, mammaglobin (US patent 5668267) or those disclosed in WO/00 52165, WO99/33869, WO99/19479, WO 98/45328. Her 2 neu antigens are disclosed inter alia, in US patent 5,801,005. Preferably the Her 2 neu comprises the entire extracellular domain ( comprising approximately amino acid 1 -  
15 645) or fragments thereof and at least an immunogenic portion of or the entire intracellular domain approximately the C terminal 580 amino acids . In particular, the intracellular portion should comprise the phosphorylation domain or fragments thereof. Such constructs are disclosed in WO00/44899. A particularly preferred construct is known as ECD PD a second is known as ECD  $\square$ PD. (See WO/00/44899.)

20 The her 2 neu as used herein can be derived from rat, mouse or human .

The plasmid may encode antigens associated with tumour-support mechanisms (e.g. angiogenesis, tumour invasion) for example tie 2, VEGF.

Vaccines of the present invention may also be used for the prophylaxis or therapy of chronic disorders in addition to allergy, cancer or infectious diseases. Such  
25 chronic disorders are diseases such as asthma, atherosclerosis, and Alzheimers and other auto-immune disorders. Vaccines for use as a contraceptive may also be considered.

Antigens relevant for the prophylaxis and the therapy of patients susceptible to or suffering from Alzheimer neurodegenerative disease are, in particular, the N  
30 terminal 39-43 amino acid fragment (A $\square$ the amyloid precursor protein and smaller fragments. This antigen is disclosed in the International Patent Application No. WO 99/27944 - (Athena Neurosciences).

Potential self-antigens that could be included as vaccines for auto-immune disorders or as a contraceptive vaccine include: cytokines, hormones, growth factors or extracellular proteins, more preferably a 4-helical cytokine, most preferably IL13. Cytokines include, for example, IL1, IL2, IL3, IL4, IL5, IL6, IL7, IL8, IL9, IL10, 5 IL11, IL12, IL13, IL14, IL15, IL16, IL17, IL18, IL20, IL21, TNF, TGF, GMCSF, MCSF and OSM. 4-helical cytokines include IL2, IL3, IL4, IL5, IL13, GMCSF and MCSF. Hormones include, for example, luteinising hormone (LH), follicle stimulating hormone (FSH), chorionic gonadotropin (CG), VGF, GHrelin, agouti, agouti related protein and neuropeptide Y. Growth factors include, for example, 10 VEGF.

The vaccines of the present invention are particularly suited for the immunotherapeutic treatment of diseases, such as chronic conditions and cancers, but also for the therapy of persistent infections. Accordingly the vaccines of the present invention are particularly suitable for the immunotherapy of infectious diseases, such 15 as Tuberculosis (TB), AIDS and Hepatitis B (HepB) virus infections.

Accordingly there is provided vaccines comprising the present invention for the immunotherapy of infectious diseases such as TB, AIDS and HepB; and their use in the manufacture of medicaments for the immunotherapy of infectious diseases such as TB, AIDS and HepB. In the context of TB, there is provided a method of treating 20 an individual suffering from TB infection, comprising the administration of a vaccine of the present invention to the individual, thereby reducing the bacterial load of that individual. The reduction of bacterial load, consisting of a reduction of the amount of TB found in the lung sputum, leading to the amelioration or cure of the TB disease.

Also, in the context of AIDS, there is provided a method of treatment of an 25 individual susceptible to or suffering from AIDS. The method comprising the administration of a vaccine of the present invention to the individual, thereby reducing the amount of CD4+ T-cell decline caused by subsequent HIV infection, or slowing or halting the CD4+ T-cell decline in an individual already infected with HIV.

30 Additionally, in the context of persistent Hepatitis B virus infection, there is provided a method of treatment of an individual susceptible to or suffering from HepB infection. Accordingly, there is provided a method comprising the administration of a vaccine of the present invention to the individual, thereby



reducing the level of HepB load in the serum (as measured by DNA clearance) and also reducing the amount of liver damage (as detected by the reduction or stabilisation of serum levels of the enzyme Alanine Transferase (ALT)).

5 The LNA-CpG/DNA complex may thus be formulated into a pharmaceutical or immunogenic composition or vaccine. In an embodiment of the invention, a polynucleotide is administered/delivered as "naked" DNA, for example as described in Ulmer et al., *Science* 259:1745-1749, 1993 and reviewed by Cohen, *Science* 259:1691-1692, 1993. Here the DNA is formulated in a buffered saline solution and injected directly into tissue. The uptake of naked DNA may be increased by coating  
10 the DNA onto biodegradable beads, which are efficiently transported into the cells or by using other well known transfection facilitating agents. LNA-conjugate/DNA may be administered in conjunction with a carrier such as, for example, liposomes. Typically such liposomes are cationic, for example imidazolium derivatives (WO95/14380), guanidine derivatives (WO95/14381), phosphatidyl choline  
15 derivatives (WO95/35301), piperazine derivatives (WO95/14651) and biguanide derivatives. The LNA-CpG/DNA complex may deliver a gene of interest such as CTFR or erythropoietin gene operatively linked to a promoter sequence. Thus a method of correcting or compensating for a disease or disorder whose etiology is characterised by a genetic aberration (such as cystic fibrosis) is provided, which  
20 method comprises the step of administering to a mammalian patient in clinical need thereof a therapeutically effective amount of the construct, preferably incorporated into a carrier.

In still another embodiment, a composition of the present invention can be delivered via a particle bombardment approach, many of which have been described  
25 (WO 91/07487). In one illustrative example, gas-driven particle acceleration can be achieved with devices such as those manufactured by Powderject Pharmaceuticals PLC (Oxford, UK) and Powderject Vaccines Inc. (Madison, WI), some examples of which are described in U.S. Patent Nos. 5,846,796; 6,010,478; 5,865,796; 5,584,807; and EP Patent No. 0500 799. This approach offers a needle-free delivery approach  
30 wherein a dry powder formulation of microscopic particles, such as polynucleotide, are accelerated to high speed within a helium gas jet generated by a hand held device, propelling the particles into a target tissue of interest, typically the skin. The particles are preferably gold beads of a 0.4 – 4.0  $\mu\text{m}$ , more preferably 0.6 – 2.0  $\mu\text{m}$  diameter

and the DNA conjugate coated onto these and then encased in a cartridge for placing into the "gene gun".

Accordingly, there is provided a DNA delivery device comprising dense microbeads coated with DNA plasmid encoding a gene of interest, which plasmid is associated with one or more LNA linked to CpG compositions of the present invention. Preferably there is provided a vaccine or immunogenic composition comprising CpG-LNA-plasmid adsorbed gold microbeads.

In a related embodiment, other devices and methods that may be useful for gas-driven needle-less injection of compositions of the present invention include those provided by Bioject, Inc. (Portland, OR), some examples of which are described in U.S. Patent Nos. 4,790,824; 5,064,413; 5,312,335; 5,383,851; 5,399,163; 5,520,639 and 5,993,412.

The invention is illustrated by, but not limited to, the following examples.

**Example 1, Investigation of LNA oligonucleotide sequence requirements and conditions for binding to supercoiled plasmid DNA**  
*Plasmids.*

Supercoiled plasmids that can be used in this study are plasmid pGG2XGFP a GFP expression vector, (Gene Therapy Systems [GTS], Inc., San Diego, California, USA), this contains multiple AG motifs within the DNA sequence of the bGHpyA region of the plasmid and multiple AAGG motifs within the DNA sequence 5' to the CMV promoter, (GTS Catalogue 2002, 50).

Plasmid pGG2XEMPTY is an expression vector, derived from pGG2XGFP by deletion of the GFP gene, but retaining a polylinker for the insertion of a gene of interest, to be expressed under the control of the CMV promoter. To construct plasmid pGG2XEMPTY from plasmid pGG2XGFP, the latter was digested with the restriction enzymes Nhe I and BamH I, ( New England Biolabs, [NEB], Hitchin, Herts., UK), to delete the region encoding the GFP gene and the remaining 5.1kb plasmid fragment was gel purified, treated with Klenow DNA Polymerase, (NEB), and ligated together using T4 DNA ligase, (NEB), prior introduction in to *E. coli*, (51). Bacterial cells containing plasmid pGG2XEMPTY were identified by standard procedures, (51 ).

*LNA oligonucleotides used in this study*

The LNA oligonucleotides used in this study are described in Table 1

5 **Table 1** lists oligonucleotide sequences used in this study.

Name	No. of sites	Description	Sequence
5876	6	13mer 100% LNA	5'-NH <sub>2</sub> - <b>CTCTCTCTCTCTC</b> -3'
5877	5	14mer 100% LNA	5'-NH <sub>2</sub> - <b>CCTTCCTTCCTTCC</b> -3'
5827	6	13mer 100% LNA	5'-NH <sub>2</sub> - <b>GAGAGAGAGAGAG</b> -3'
5875	6	11mer 100% LNA	5'-NH <sub>2</sub> - <b>CTCTCTCTCTC</b> -3'
5747	8	9mer 'bis' 50% LNA	5'-NH <sub>2</sub> - <b>CtCtCtCtC-XXX- CtCtCtCtC</b> -3'
6563	6	11mer 100% LNA	5'-TAMRA- <b>CTCTCTCTCTC</b> -3'
11701	5	14mer 100% LNA	5'-NH <sub>2</sub> - <b>GGAAGGAAGGAAGG</b> -3'
PTOCpG	6	21mer DNA13mer LNA	5' <i>tccatgacgttcctgacgttt</i> <b>GAGAGAGAGA</b> <b>GAG-3'</b>
PTOGpC	6	21mer DNA13mer LNA	5' <i>tccatgagcttcctgagtcct</i> <b>GAGAGAGAGA</b> <b>GAG-3'</b>
5'SHGA	6	13mer 100% LNA	5'-S-S- <b>GAGAGAGAGAGAG</b> -3'

LNA residues are displayed in bold upper case, DNA residues are shown in bold lower case with PTO residues additionally italicised. Number of sites refers to the maximum number of theoretical oligonucleotide binding sites present on either the

10 gWiz or pGG2XGFP, plasmid.

X = 'PEG spacer' - 9-O-Dimethoxytrityl-triethylene glycol, 1-[(2-cyanoethyl)-(N,N-diisopropyl)]-phosphoramidite, spacer phosphoramidate 9, (Glen Research, USA); O = 8-amino-3,6-dioxaoctanoic acid linker, g = glycine, l = lysine, Fl = Fluorescein, NH<sub>2</sub> = 5'- amino-modifier C12 phosphoramidite spacers, (Glen Research, USA), PTO = phosphorothioate, S-S = Thiol modifier, C6 S-S phosphoramidate, (Glen Research, USA).

15

PG5044

All LNA oligonucleotides were synthesized by Proligo LLC, Colorado, USA. The majority were made with 5'- amino-modifier C12 phosphoramidite spacers, (Glen Research, USA), to allow for labelling with Alexa Fluor dyes, (Molecular Probes, Netherlands), or heterobifunctional linkers, eg. Maleimide or SPDP, (Perbio, USA).

5 Most are 100% LNA monomers, but LNA 5747, (Table 3), is 50% LNA and 50% DNA. This 'bis' LNA oligonucleotide was made as an analogue of the 'bis' PNA clamps described in Example 1, (50, 52), and could only be efficiently synthesized as a 50:50 mix of LNA and DNA residues.

#### 10 *Analysis of conditions for binding of LNA oligonucleotides to supercoiled plasmid DNA*

Annealing / labelling conditions for LNA oligonucleotides were based upon those described in the literature, (55). In order to maximise the efficiency of PNA binding to supercoiled plasmid DNA labelling was performed in a buffer containing  
15 no salt at low pH < 6, ( 10mM phosphate buffer, 1mM EDTA, pH 7) for 16 hours at 37°C. The low pH should enable cytosine residues to Hoogsten base pair with similar efficiency to pseudocytosine residues, (52), Initially, 10ug of plasmid DNA was labelled in a total volume of 20ul, where PNA oligonucleotides were present at a 20 X molar excess over the maximum number of potential binding sites present in the  
20 plasmid DNA, 10 sites, ( see Table 1, 50). For OsPNA2, labelling was performed by heating at 95°C for 10 minutes followed by 10 minutes at room temperature, (20°C), following by 16 hours at 37°C, (7).

For the conditions described above LNAs could be detected as bound to plasmid.

25

#### **OTHER PUBLICATIONS.**

- 1) Rajur *et al.*, 8(6):935-940 (1997).
- 2) Brinkley, M., 3(1):2-13 (1992).
- 30 3) Mattaj, I & Engelmeier, L., Annual Review Biochemistry.67:265-306 (1998)
- 4) Gorlich, D. & Kutay, U., Annual Review Cellular & Developmental Biology 15:607-660 (1999).
- 5) Zanta et al., Proc. Natl. Acad. Sci. USA 96: 91-96 (1999).

- 6) Sebestyen et al., *Nature Biotech.* 16: 80-85 (1998).
- 7) Branden et al., *Nature* 17: 784-787 (1999).
- 8) Zhang et al., *Gene Therapy* 6: 171-181 (1999).
- 9) Subramanian et al., *Nature* 17: 873-877 (1999).
- 5 10) Chang et al., *Virology* 189: 821-827 (1992).
- 11) Vives et al., *Journal Biological Chemistry* 272 (25): 16010-16017 (1997).
- 12) Derossi et al., *Journal Biological Chemistry* 269 (14):10444-10450 (1994).
- 13) Derossi et al., *Trends in Cell Biology* 8: 84-87 (1998).
- 14) Prochanitz, A., *Current Opinion in Neurobiology* 6: 629-634 (1996).
- 10 15) Prochiantz, A. & Theodore, L., *BioEssays* 17(1): 39-44 (1995).
- 16) Eguchi et al., *J. Biol Chem* 272: 17640-17647 (1997).
- 17) Burglin, T. & De Robertis, E., *EMBO J.* 6: 2617-2625 (1987).
- 18) Ballas, N. & Citovsky, V., *Proc. Natl. Acad. Sci. USA* 94: 10723-10728 (1997).
- 19) Nigg, E., *Nature* 386: 779-787 (1997).
- 15 20) Blanke et al., *Proc. Natl. Acad. Sci. USA* 93: 8437-8442 (1996).
- 21) Morris et al., *Nucleic Acids Research* 27 3510-3517 (1999).
- 22) Schmolke et al., *Journal of Virology* 69: 1071-1078 (1995).
- 23) Wychowski et al., *EMBO J.* 5: 2569-2576.
- 24) Xia et al., *Journal of Virology* 66: 914-921 (1992).
- 20 25) Dean, D., *Adv. Drug Delivery reviews* 44: 81-95 (2000).
- 26) Lindgren et al., *Trends in Pharmaceutical Sciences* 21: 99-103 (2000).
- 27) Elliot, G. & O' Hare, P, *Cell* 88: 223-233 (1997).
- 28) Kawana et al., *Journal of Virology* 75 (5): 2331-2336 (2001).
- 29) Pastan, I & Fitzgerald, D., *Science* 254: 1173-1177 (1991).
- 25 30) Oess, S. & Hildt, E., *Gene Therapy* 7: 750-758 (2000).
- 31) Mechtler, K. & Wagner, E, *New Journal of Chemistry* 21: 105-111 (1997).
- 32) Pillot et al., *Journal Biological Chemistry* 271: 28757-28765 (1996).
- 33) Subbarao et al., *Biochemistry* 26 (11) : 2964-2972 (1987).
- 34) Niidome et al., *Journal Biological Chemistry*, 272: 15307-15312 (1997).
- 30 35) Wyman et al., *Biochemistry* 36: 3008-3017 (1997).
- 36) Vogel et al., *Journal American Chemical Society* 118: 1581-1586 (1996).
- 37) Kichler et al., *Bioconjugate Chemistry* 8: 213-221 (1997).
- 38) Puls, R. & Minchin, M., *Gene Therapy* 6: 1774-1778 (1999).

- 39) Goldman et al., *Cancer Research* 57: 1447-1451 (1997)
- 40) Kircheis et al., *Gene Therapy* 4: 409-418 (1997)
- 41) Kawakami et al., *Gene Therapy* 7: 292-299 (2000).
- 42) Fajac et al., *Human Gene Therapy* 10: 395-406 (1999).
- 5 43) Zanta et al., *Bioconjugate Chemistry* 8: 839-844 (1997).
- 44) Hung et al., *Cancer Research* 61: 1080-1088 (2001).
- 45) Hart et al., *Gene Therapy*, 2: 552-554 (1995).
- 46) Deonarain, M., *Expert Opinion on Therapeutic Patents* 8(1): 53-69 (1998).
- 47) Cristiano, R. & Roth, J., *Journal Molecular Medicine* 73: 479-486 (1995)
- 10 48) Uherek, C. et al., *Journal of Biological Chemistry*, 273: 8835-8841 (1998).
- 49) Chattergoon et al., *Nature Biotechnology* 18: 974-979 (2000).
- 50) Zelphati et al., *Human Gene Therapy* 10: 15-24 (1999).
- 51) Maniatis et al., 'Molecular Cloning: a laboratory manual' 2<sup>nd</sup> Edition, (1992)
- 52) Egholm et al., *Nucleic Acids Research* 23: 217-222 (1995).
- 15 53) Zhang et al., *Nucleic Acids Research* 28 (17): 3332- 3338 (2000).
- 54) Griffith et al., *Journal American Chemical Society* 117: 831-832, (1995).
- 55) Nielsen, P. & Egholm, M. 'An introduction to PNA' Chapter 1 In Nielsen, P. & Egholm, M (Eds.) *Peptide Nucleic Acids: Protocols and Applications* (1999).
- 56) Butz, E. & Bevan, M., *Journal of Immunology* 160: 2139-2144, (1998).
- 20 57) Tuting, et al., *Journal of Immunology* 160: 1139-1147, (1998).
- 58) Harrison, et al., *Vaccine* 19: 1820-1826, (2001).
- 59) Vasilakos, et al., *Cellular immunology* 204: 64-74 (2000).
- 60) Rhodes, J. et al., *Nature* 377: 71-75 (1995).
- 61) Gherardi, M. et al., *Journal of Virology* 74 (14): 6278-6286.
- 25 62) Alderton, W. et al., *Biochemical Journal* 357: 593-615, (2001).
- 63) Scheerlinck, JP. 'Genetic adjuvants for DNA vaccines', *Vaccine* 19: 2647-2656, (2001).
- 64) Reyes et al., *Vaccine* 19: 3778-3786, (2001).
- 65) Sato, Y. et al., *Science* 273 (5273): 352-354 (1996).
- 30 66) Hemmi, H. et al., *Nature* 408: 740-745, (2000)
- 67) Beutler, B., *Current Opinion in Microbiology* 3: 23-30 (2000).
- 68) Thoma-Uszynski, S. et al., *Journal of Immunology* 165: 3804-3810 (2000).
- 69) Mitchell, P. & Tijan, R., *Science* 245: 371-378, (1989).

- 70) Courey, A. & Tijan, R., *Cell* 55: 887-898, (1988).
- 71) Tanaka et al., *Molecular and Cellular Biology* 14: : 6046-6055, (1994).
- 72) Ma, J. & Ptashne, M., *Cell*, 51: 113-119, (1987).
- 73) Seipel et al., *EMBO Journal* 11: 4961-4968, (1992).
- 5 74) Obika et al., *Tetrahedron Letters* 41: 221-224, (2000).
- 75) Torigue et al., *Journal of Biological Chemistry* 276(4): 2354-2360, (2001).
- 76) Smulevitch et al., *Nature Biotechnology* 14: 1700-1704, (1996).
- 77) Kichler et al., *Bioconjugate Chemistry* 8: 213-221, (1997).
- 78) Bukanov et al., *Proc. Natl. Acad. Sci. USA* 95: 5516-5518, (1998).
- 10 79) Klinman et al., *Vaccine* 17: 19-25, (1999).
- 80) Weeratna et al., *Vaccine* 18: 1755-1762, (2000).
- 81) Shirota et al., *Journal of Immunology* 167: 66-74, (2001).
- 82) Hacker et al., *EMBO Journal* 17: 6230-6240, (1998).
- 83) Bauer et al., *Proc. Natl. Acad. Sci. USA* 98 (16): 9237-9242, (2001); International  
15 patent application: ' Process for high throughput screening of CpG-based immuno-  
agonist/antagonists', WO02/228009A2, (2002); S. Bauer, personal communication,  
(unpublished observation).
- 84) Jin et al., *Journal of Immunology* 165: 5153-5160, (2000).
- 85) Dalpke et al., *Journal of Immunology* 166: 7082-7089, (2001).
- 20 86) Crabtree et al., *Infection and Immunity* 69(4): 2123-2129, (2001).
- 87) Gao et al., *Journal of Immunology* 166: 6855-6860, (2001).
- 88) Tung, C.H. and Stein, S., *Bioconjugate Chemistry* 11(5): 605-618, (2000).
- 89) Wang et al., *Bioconjugate Chemistry* 8: 878-884, (1997).
- 90) Harrison, J. and Balasubramanian, S., *Nucleic Acids Research* 26 (13): 3136-  
25 3145, (1998).
- 91) Stetsenko, D. and Gait, M., *J. Org. Chem.* 65: 4900-4908, (2000).
- 92) Stetsenko, D. and Gait, M., *Nucleosides, Nucleotides & Nucleic Acids* 19(10-12):  
1751-1764, (2000).
- 93) Stetsenko, D. and Gait, M., *Bioconjugate Chemistry* 12: 576-586, (2001).
- 30 94) Rojas et al., *Nature Biotechnology* 16: 370-375, (1998).
- 95) Ogris et al., *Journal of Biochemistry* 276 (50): 47550-47555, (2001).

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☒ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**